

=> s transaldolase/cn  
L1 1 TRANSALDOLASE/CN

=> d

L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS

RN 9014-46-4 REGISTRY

CN **Transaldolase (9CI)** (CA INDEX NAME)

OTHER NAMES:

CN Dihydroxyacetone transferase

CN E.C. 2.2.1.2

MF Unspecified

CI MAN

LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA, CAPLUS,  
CHEMCATS, CHEMLIST, EMBASE, TOXCENTER, USPAT2, USPATFULL

Other Sources: EINECS\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

437 REFERENCES IN FILE CA (1957 TO DATE)

2 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

438 REFERENCES IN FILE CAPLUS (1957 TO DATE)

=> d his'

(FILE 'HOME' ENTERED AT 15:23:58 ON 08 JUL 2003)

L1 FILE 'REGISTRY' ENTERED AT 15:24:05 ON 08 JUL 2003  
1 S TRANSALDOLASE/CN

FILE 'HCAPLUS' ENTERED AT 15:25:27 ON 08 JUL 2003

L2 FILE 'REGISTRY' ENTERED AT 15:25:32 ON 08 JUL 2003  
SET SMARTSELECT ON  
SEL L1 1- CHEM : 4 TERMS  
SET SMARTSELECT OFF

L3 FILE 'HCAPLUS' ENTERED AT 15:25:33 ON 08 JUL 2003  
664 S L2  
L4 4 S L3 (L) (CORYNEBACTERIA OR CORYNEBACTERIA GLUTAMICUM OR (BACTE

=> d ibib ab 1-4

L4 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 2001:246415 HCAPLUS  
DOCUMENT NUMBER: 135:328890  
TITLE: Modeling and Experimental Design for Metabolic Flux  
Analysis of Lysine-Producing Corynebacteria by Mass  
Spectrometry  
AUTHOR(S): Wittmann, Christoph; Heinzle, Elmar  
CORPORATE SOURCE: Biochemical Engineering Institute, Saarland  
University, Saarbruecken, Germany  
SOURCE: Metabolic Engineering (2001), 3(2), 173-191  
CODEN: MEENFM; ISSN: 1096-7176  
PUBLISHER: Academic Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Exptl. design of <sup>13</sup>C-tracer studies for metabolic flux anal. with mass spectrometric detn. of labeling patterns was performed for the central metab. of Corynebacterium glutamicum comprising various flux scenarios. Ratio measurement of mass isotopomer pools of Corynebacterium products lysine, alanine, and trehalose is sufficient to quantify the flux partitioning ratios (i) between glycolysis and pentose phosphate pathways (.PHI.PPP), (ii) between the split pathways in the lysine biosynthesis (.PHI.DH), (iii) at the pyruvate node (.PHI.PC), and the reversibility of (iv) glucose 6-phosphate isomerase (.zeta.PGI), (v) at the pyruvate node (.zeta.PC/PEPCK), and (vi) of transaldolase and transketolases in the PPP. Weighted sensitivities for flux parameters were derived from partial derivs. to quant. evaluate exptl. approaches and predict precision for estd. flux parameters. Deviation of intensity ratios from ideal values of 1 was used as weighting function. Weighted flux sensitivities can be used to identify optimal type and degree of tracer labeling or potential intensity ratios to be measured. Exptl. design for lysine-producing strain C. glutamicum MH 20-22B and various potential mutants with different alterations in the flux pattern showed that specific tracer labeling is optimal to quantify a certain flux parameter uninfluenced by the overall flux situation. Identified substrates of choice are [1-<sup>13</sup>C]glucose for the estn. of .PHI.PPP and .zeta.PGI and a 1:1 mixt. of [U-<sup>12</sup>C/U-<sup>13</sup>C]glucose for the detn. of .zeta.PC/PEPCK. .PHI.PC can be quantified by feeding [4-<sup>13</sup>C]glucose or [U-<sup>12</sup>C/U-<sup>13</sup>C]glucose (1:1), whereas .PHI.DH is accessible via [4-<sup>13</sup>C]glucose. The sensitivity for the quantification of a certain flux parameter can be influenced by superposition through other flux parameters in the network, but substrate and measured mass isotopomers of choice remain the same. In special cases, reduced labeling degree of the tracer substrate can increase the precision of flux anal. Enhanced precision and flux information can be achieved via multiply labeled substrates. The presented approach can be applied for effective exptl. design of <sup>13</sup>C tracer studies for metabolic flux anal. Intensity ratios of other products such as glutamate, valine, phenylalanine, and riboflavin also sensitively reflect flux parameters, which underlines the great potential of mass spectrometry for flux anal.  
(c) 2001 Academic Press.

REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 2001:50828 HCAPLUS  
DOCUMENT NUMBER: 134:111274  
TITLE: Sequences of Coryneform bacteria tal gene and uses  
thereof in fermentative preparation of L-amino acids  
INVENTOR(S): Dunican, L. K.; McCormack, Ashling; Stapelton, Cliona;  
Burke, Kevin; Mockel, Bettina  
PATENT ASSIGNEE(S): Degussa-Huls Aktiengesellschaft, Germany; National  
University of Ireland  
SOURCE: PCT Int. Appl., 47 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001004325	A1	20010118	WO 2000-EP6304	20000705
W: AU, BR, CA, CN, HU, ID, JP, KR, MX, PL, RU, SK, UA, ZA				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP 1109915	A1	20010627	EP 2000-956165	20000705
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 2000006915	A	20010731	BR 2000-6915	20000705
PRIORITY APPLN. INFO.:				
			US 1999-142915P	P 19990709
			US 2000-531266	A 20000320
			WO 2000-EP6304	W 20000705

AB The invention provides protein and DNA sequences of tal genes from coryneform bacteria. The invention further provides new measures for improved fermentative prepn. of amino acids, in particular L-lysine, L-threonine, L-isoleucine and L-tryptophan.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 2001:50825 HCAPLUS

DOCUMENT NUMBER: 134:111273

TITLE: Sequences of Coryneform bacteria opcA gene and uses thereof in fermentative preparation of L-amino acids

INVENTOR(S): Dunican, L. K.; McCormack, Ashling; Stapelton, Cliona; Burke, Kevin; Moritz, Bernd; Eggeling, Lothar; Sahm, Hermann; Mockel, Bettina; Weissenborn, Anke

PATENT ASSIGNEE(S): Degussa-Huls Aktiengesellschaft, Germany; Forschungszentrum Julich G.m.b.H.; National University of Ireland

SOURCE: PCT Int. Appl., 75 pp.  
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001004322	A1	20010118	WO 2000-EP6300	20000705
WO 2001004322	C2	20020912		
W: AU, BR, CA, CN, HU, ID, JP, KR, MX, PL, RU, SK, UA, ZA				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
BR 2000006909	A	20010612	BR 2000-6909	20000705
EP 1109913	A1	20010627	EP 2000-945874	20000705
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
PRIORITY APPLN. INFO.:				
			US 1999-142915P	P 19990709
			US 2000-531267	A 20000320
			WO 2000-EP6300	W 20000705

AB The invention provides protein and DNA sequences of opcA genes from coryneform bacteria. The invention further provides new measures for improved fermentative prepn. of amino acids, in particular L-lysine, L-threonine, L-isoleucine and L-tryptophan.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 1962:26672 HCAPLUS

DOCUMENT NUMBER: 56:26672

ORIGINAL REFERENCE NO.: 56:5110h-i

TITLE: Pentose cycle in Corynebacterium diphtheriae

AUTHOR(S): Halanicka, Danuta

CORPORATE SOURCE: Inst. Mother Child, Warsaw, Polish

SOURCE: Acta Biochimica Polonica (1960), 7, 449-57

DOCUMENT TYPE:

Journal

LANGUAGE:

Unavailable

AB Enzymes of the pentose cycle in Me<sub>2</sub>CO powder exts. and in the supernatants (18,500 g, 30 min.) of *Corynebacterium diphtheriae* were investigated. The preps. metabolized ribose 5-phosphate and produced keto sugars. Identification of ribose 5-phosphate was performed spectrophotometrically by means of the Dische reaction. Formation of sedoheptulose and its transition into hexose was shown. The presence of the following enzymes in *Corynebacteria* was found: phosphopentose isomerase, epimerase, transketolase, **transaldolase**, glucose-6-phosphate dehydrogenase, 6-phosphogluconic acid dehydrogenase, and ribokinase. Dehydrogenases were linked with triphosphopyridine nucleotide, the kinase was specific for ribose.

RESULT 4

AAF25333

ID AAF25333 standard; DNA; 1083 BP.

XX

AC AAF25333;

XX

DT 30-APR-2001 (first entry)

XX

DE Coding region of the tal gene.

XX

KW tal gene; Coryneform bacteria; L-amino acid; transaldolase; ss.

XX

OS Corynebacterium glutamicum.

XX

FH Key Location/Qualifiers

FT CDS 1..1083

FT /\*tag= a

FT /product= "tal"

XX

PN WO200104325-A1.

XX

PD 18-JAN-2001.

XX

PF 05-JUL-2000; 2000WO-EP06304.

XX

PR 09-JUL-1999; 99US-0142915.

PR 20-MAR-2000; 2000US-0531266.

XX

PA (DEGS ) DEGUSSA-HUELS AG.

PA (UYNA-) UNIV NAT IRELAND.

XX

PI Dunican LK, McCormack A, Stapelton C, Burke K, Moeckel B;

XX

DR WPI; 2001-159407/16.

DR P-PSDB; AAB31783.

XX

PT New polynucleotides from coryneform bacteria, specifically

PT Corynebacterium, useful for preparing L-amino acids, especially

PT L-lysine, L-threonine, L-isoleucine and L-tryptophan, by amplifying tal

PT gene -

XX

PS Claim 4; Page 43-45; 48pp; English.

XX

CC The present sequence represents a tal gene fragment from Coryneform  
 CC bacteria. Tal polynucleotides and polypeptides are used for fermentative  
 CC preparation of L-amino acids, especially L-lysine, L-threonine,  
 CC L-isoleucine and/or L-tryptophan. The tal polynucleotide is useful as  
 CC a hybridisation probe for isolating a cDNA encoding for the tal gene  
 CC product, and for isolating cDNA or genes having high similarity with  
 CC the sequence of the tal gene. The polynucleotides may be used as  
 CC hybridisation probes for RNA, cDNA and DNA to isolate full-length cDNA  
 CC which code for transaldolase and to isolate those cDNA or genes, which  
 CC have a high similarity with that of the transaldolase gene. These may  
 CC also be used as primers for the preparation of DNA which code for  
 CC transaldolase by polymerase chain reaction (PCR).

XX

SQ Sequence 1083 BP; 215 A; 336 C; 294 G; 238 T; 0 other;

Query Match 98.7%; Score 1065.6; DB 22; Length 1083;  
Best Local Similarity 99.2%; Pred. No. 1e-269;  
Matches 1071; Conservative 0; Mismatches 9; Indels 0; Gaps 0;

Qy	1	ATGTCTCACATTGATGATCTTGACAGCTCGGCACCTTCCACTTGGCTCGACGACCTCTCC	60
Db	1	ATGTCTCACATTGATGATCTTGACAGCTCGGCACCTTCCACTTGGCTCGACGACCTCTCC	60
Qy	61	CGCGAGCGCATTACTTCCGGCAATCTCAGCCAGGTTATTGAGGAAAAGTCTGTAGTCGGT	120
Db	61	CGCGAGCGCATTACTTCCGGCAATCTCAGCCAGGTTATTGAGGAAAAGTCTGTAGTCGGT	120
Qy	121	GTCACCACCAACCCAGCTATTTTCGCAGCAGCAATGTCCAAGGGCGATTCCCTACGACGCT	180
Db	121	GTCACCACCAACCCAGCTATTTTCGCAGCAGCAATGTCCAAGGGCGATTCCCTACGACGCT	180
Qy	181	CAGATCGCAGAGCTCAAGGCCGCTGGCGCATCTGTTGACCAGGCTGTTTACGCCATGAGC	240
Db	181	CAGATCGCAGAGCTCAAGGCCGCTGGCGCATCTGTTGACCAGGCTGTTTACGCCATGAGC	240
Qy	241	ATCGACGATGTTTCGCAATGCTTGTGATCTGTTACCGGCATCTTCGAGTCCTCCAACGGC	300
Db	241	ATCGACGATGTTTCGCAATGCTTGTGATCTGTTACCGGCATCTTCGAGTCCTCCAACGGC	300
Qy	301	TACGACGGCCGCGTGTCCATCGAGGTTGACCCACGTATCTCTGCTGACCGCGACGCAACC	360
Db	301	TACGACGGCCGCGTGTCCATCGAGGTTGACCCACGTATCTCTGCTGACCGCGACGCAACC	360
Qy	361	CTGGCTCAGGCCAAGGAGCTGTGGGCAAAGGTTGATCGTCCAAACGTCATGATCAAGATC	420
Db	361	CTGGCTCAGGCCAAGGAGCTGTGGGCAAAGGTTGATCGTCCAAACGTCATGATCAAGATC	420
Qy	421	CCTGCAACCCCAGGTTCTTTGCCAGCAATCACCGACGCTTTGGCTGAGGGCATCAGCGTT	480
Db	421	CCTGCAACCCCAGGTTCTTTGCCAGCAATCACCGACGCTTTGGCTGAGGGCATCAGCGTT	480
Qy	481	AACGTACCTTGATCTTCTCCGTTGCTCGCTACCGCGAGGTCATCGCTGCGTACATCGAG	540
Db	481	AACGTACCTTGATCTTCTCCGTTGCTCGCTACCGCGAGGTCATCGCTGCGTTCATCGAG	540
Qy	541	GGAATCAAGCAGGCAGCTGCAAACGGCCACGACGTATCCAAGATCCACTCTGTGGCTTCC	600
Db	541	GGCATCAAGCAGGCTGCTGCAAACGGCCACGACGTCTCCAAGATCCACTCTGTGGCTTCC	600
Qy	601	TTCTTCGTCTCCCGCGTCGACGTTGAGATCGACAAGCGCCTCGAGGCAATCGGATCCGAT	660
Db	601	TTCTTCGTCTCCCGCGTCGACGTTGAGATCGACAAGCGCCTCGAGGCAATCGGATCCGAT	660
Qy	661	GAGGCTTTGGCTCTGCGCGGCAAGGCAGGCGTTGCCAACGCTCAGCGCGCTTACGCTGTG	720
Db	661	GAGGCTTTGGCTCTGCGCGGCAAGGCAGGCGTTGCCAACGCTCAGCGCGCTTACGCTGTG	720
Qy	721	TACAAGGAGCTTTTCGACGCCGCCGAGCTGCCTGAAGGTGCCAACACTCAGCGCCCACTG	780
Db	721	TACAAGGAGCTTTTCGACGCCGCCGAGCTGCCTGAAGGTGCCAACACTCAGCGCCCACTG	780

[illegible]



RESULT 4

AAF25333

ID AAF25333 standard; DNA; 1083 BP.

XX

AC AAF25333;

XX

DT 30-APR-2001 (first entry)

XX

DE Coding region of the tal gene.

XX

KW tal gene; Coryneform bacteria; L-amino acid; transaldolase; ss.

XX

OS Corynebacterium glutamicum.

XX

FH Key Location/Qualifiers

FT CDS 1..1083

FT /\*tag= a

FT /product= "tal"

XX

PN WO200104325-A1.

XX

PD 18-JAN-2001.

XX

PF 05-JUL-2000; 2000WO-EP06304.

XX

PR 09-JUL-1999; 99US-0142915.

PR 20-MAR-2000; 2000US-0531266.

XX

PA (DEGS ) DEGUSSA-HUELS AG.

PA (UYNA-) UNIV NAT IRELAND.

XX

PI Dunican LK, McCormack A, Stapelton C, Burke K, Moeckel B;

XX

DR WPI; 2001-159407/16.

DR P-PSDB; AAB31783.

XX

PT New polynucleotides from coryneform bacteria, specifically

PT Corynebacterium, useful for preparing L-amino acids, especially

PT L-lysine, L-threonine, L-isoleucine and L-tryptophan, by amplifying tal

PT gene -

XX

PS Claim 4; Page 43-45; 48pp; English.

XX

CC The present sequence represents a tal gene fragment from Coryneform  
 CC bacteria. Tal polynucleotides and polypeptides are used for fermentative  
 CC preparation of L-amino acids, especially L-lysine, L-threonine,  
 CC L-isoleucine and/or L-tryptophan. The tal polynucleotide is useful as  
 CC a hybridisation probe for isolating a cDNA encoding for the tal gene  
 CC product, and for isolating cDNA or genes having high similarity with  
 CC the sequence of the tal gene. The polynucleotides may be used as  
 CC hybridisation probes for RNA, cDNA and DNA to isolate full-length cDNA  
 CC which code for transaldolase and to isolate those cDNA or genes, which  
 CC have a high similarity with that of the transaldolase gene. These may  
 CC also be used as primers for the preparation of DNA which code for  
 CC transaldolase by polymerase chain reaction (PCR).

XX

SQ Sequence 1083 BP; 215 A; 336 C; 294 G; 238 T; 0 other;

Query Match 98.7%; Score 1065.6; DB 22; Length 1083;  
Best Local Similarity 99.2%; Pred. No. 1e-269;  
Matches 1071; Conservative 0; Mismatches 9; Indels 0; Gaps 0;

Qy	1	ATGTCTCACATTGATGATCTTGACACAGCTCGGCACCTTCCACTTGGCTCGACGACCTCTCC	60
Db	1	ATGTCTCACATTGATGATCTTGACACAGCTCGGCACCTTCCACTTGGCTCGACGACCTCTCC	60
Qy	61	CGCGAGCGCATTACTTCCGGCAATCTCAGCCAGGTTATTGAGGAAAAGTCTGTAGTCGGT	120
Db	61	CGCGAGCGCATTACTTCCGGCAATCTCAGCCAGGTTATTGAGGAAAAGTCTGTAGTCGGT	120
Qy	121	GTCACCACCAACCCAGCTATTTTCGCAGCAGCAATGTCCAAGGGCGATTCTACGACGCT	180
Db	121	GTCACCACCAACCCAGCTATTTTCGCAGCAGCAATGTCCAAGGGCGATTCTACGACGCT	180
Qy	181	CAGATCGCAGAGCTCAAGGCCGCTGGCGCATCTGTTGACCAGGCTGTTTACGCCATGAGC	240
Db	181	CAGATCGCAGAGCTCAAGGCCGCTGGCGCATCTGTTGACCAGGCTGTTTACGCCATGAGC	240
Qy	241	ATCGACGATGTTTCGCAATGCTTGATCTGTTACCGGCATCTTCGAGTCCTCCAACGGC	300
Db	241	ATCGACGACGTTTCGCAATGCTTGATCTGTTACCGGCATCTTCGAGTCCTCCAACGGC	300
Qy	301	TACGACGGCCGCGTGTCCATCGAGGTTGACCCACGTATCTCTGCTGACCGCGACGCAACC	360
Db	301	TACGACGGCCGCGTGTCCATCGAGGTTGACCCACGTATCTCTGCTGACCGCGACGCAACC	360
Qy	361	CTGGCTCAGGCCAAGGAGCTGTGGGCAAAGGTTGATCGTCCAAACGTCATGATCAAGATC	420
Db	361	CTGGCTCAGGCCAAGGAGCTGTGGGCAAAGGTTGATCGTCCAAACGTCATGATCAAGATC	420
Qy	421	CCTGCAACCCAGGTTCTTTGCCAGCAATCACCGACGCTTTGGCTGAGGGCATCAGCGTT	480
Db	421	CCTGCAACCCAGGTTCTTTGCCAGCAATCACCGACGCTTTGGCTGAGGGCATCAGCGTT	480
Qy	481	AACGTCACCTTGATCTTCTCCGTTGCTCGCTACCGCGAGGTCATCGCTGCGTTACATCGAG	540
Db	481	AACGTCACCTTGATCTTCTCCGTTGCTCGCTACCGCGAGGTCATCGCTGCGTTACATCGAG	540
Qy	541	GGAATCAAGCAGGCAGCTGCAAACGGCCACGACGTATCCAAGATCCACTCTGTGGCTTCC	600
Db	541	GGCATCAAGCAGGCTGCTGCAAACGGCCACGACGTCTCCAAGATCCACTCTGTGGCTTCC	600
Qy	601	TTCTTCGTCTCCCGCGTCGACGTTGAGATCGACAAGCGCCTCGAGGCAATCGGATCCGAT	660
Db	601	TTCTTCGTCTCCCGCGTCGACGTTGAGATCGACAAGCGCCTCGAGGCAATCGGATCCGAT	660
Qy	661	GAGGCTTTGGCTCTGCGCGGCAAGGCAGGCGTTGCCAACGCTCAGCGCGCTTACGCTGTG	720
Db	661	GAGGCTTTGGCTCTGCGCGGCAAGGCAGGCGTTGCCAACGCTCAGCGCGCTTACGCTGTG	720
Qy	721	TACAAGGAGCTTTTCGACGCCGCCGAGCTGCCTGAAGGTGCCAACACTCAGCGCCCACTG	780
Db	721	TACAAGGAGCTTTTCGACGCCGCCGAGCTGCCTGAAGGTGCCAACACTCAGCGCCCACTG	780

Qy	781	TGGGCATCCACCGGCGTGAAGAACCCTGCGTACGCTGCAACTCTTTACGTTTCCGAGCTG	840
Db	781	TGGGCATCCACCGGCGTGAAGAACCCTGCGTACGCTGCAACTCTTTACGTTTCCGAGCTG	840
Qy	841	GCTGGTCCAAACACCGTCAACACCATGCCAGAAGGCACCATCGACGCTGTTCGGAAGTG	900
Db	841	GCTGGTCCAAACACCGTCAACACCATGCCAGAAGGCACCATCGACGCGGTTCTGGAGCAG	900
Qy	901	GGCAACCTGCACGGTGACACCCTGTCCAACCTCCGCGGCAGAAGCTGACGCTGTGTTCTCC	960
Db	901	GGCAACCTGCACGGTGACACCCTGTCCAACCTCCGCGGCAGAAGCTGACGCTGTGTTCTCC	960
Qy	961	CAGCTTGAGGCTCTGGGCGTTGACTTGGCAGATGTCTTCCAGGTCCTGGAGACCGAGGGT	1020
Db	961	CAGCTTGAGGCTCTGGGCGTTGACTTGGCAGATGTCTTCCAGGTCCTGGAGACCGAGGGT	1020
Qy	1021	GTGGACAAGTTTGTGCTTCTTGGAGCGAAGTGCTTGAGTCCATGGAAGCTCGCCTGAAG	1080
Db	1021	GTGGACAAGTTTGTGCTTCTTGGAGCGAAGTGCTTGAGTCCATGGAAGCTCGCCTGAAG	1080